

U. S. AIR FORCE

PROJECT RAND

RESEARCH MEMORANDUM

AN INFINITE-DIMENSIONAL EXTENSION
OF A SYMETRIC BLOTTO GAME

Oliver Gross

RM-718

November 8, 1951

Assigned to _____

This is a working paper. It may be expanded,
modified, or withdrawn at any time.

The **RAND** *Corporation*

1700 MAIN ST. • SANTA MONICA • CALIFORNIA

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 08 NOV 1951		2. REPORT TYPE		3. DATES COVERED 00-00-1951 to 00-00-1951	
4. TITLE AND SUBTITLE An Infinite-Dimensional Extension of a Symetric Blotto Game				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Rand Corporation, Project Air Force, 1776 Main Street, PO Box 2138, Santa Monica, CA, 90407-2138				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Summary: A two-person zero-sum game over a function space and with discontinuous payoff is considered which has an optimal strategy for either player, consisting of randomizing over a certain two-parameter family of functions.

AN INFINITE-DIMENSIONAL EXTENSION OF A SYMMETRIC BLOTTO GAME

Oliver Gross

Let X be the set of all real-valued continuous functions over $[0,1]$ with the properties

$$x(t) \geq 0 \text{ for all } t \in [0,1] \text{ and}$$

$$\int_0^1 x(t) dt = 1$$

for any $x \in X$. We set up the normal form of the game thus:

1. A pure strategy for the maximizing player is a point $x \in X$.
2. A pure strategy for the minimizing player is a point $y \in X$.
3. For any selection of pure strategies $\{x, y\}$, the payoff M to the maximizing player is given by

$$M(x, y) = \int_0^1 v(t) \operatorname{sgn} [x(t) - y(t)] dt,$$

where v is a prescribed positive-valued element of X .

Remarks

One can envision this game as two lines of infantry of equal strength about to engage in battle in a corridor, the value of a breakthrough to either force in any interval dt along the width of the corridor being $v(t)dt$. A breakthrough is achieved at the interval, if "Blotto's" forces exceed the enemy's there, or conversely. We note that this game is an extension of the finite-dimensional cases considered in [1] and [2].

We propose that the following is an optimal strategy for the maximizing player (hereafter denoted by "Blotto"). It would follow by symmetry that the enemy could do equally as well by employing the same strategy. Hence, to prove we have a solution, we need only verify that the following strategy will ensure Blotto an expectation of zero:

Blotto chooses one of the following 2-parameter family of functions:

$$(1) \quad x_{\alpha, \rho}(t) = v(t) \left[1 + \rho \cos 2\pi \left(\alpha - \int_0^t v(\lambda) d\lambda \right) \right],$$

where the choice is affected by selecting α and ρ independently from the respective distributions

$$(2) \quad f_1^*(\alpha) = \alpha \quad 0 \leq \alpha \leq 1,$$

$$(3) \quad f_2^*(\rho) = 1 - \sqrt{1 - \rho^2} \quad 0 \leq \rho \leq 1.$$

Verification

We first show that the foregoing is admissible as a mixed strategy. It should suffice to show that (i) f_1^* and f_2^* are distributions over $[0,1]$, and (ii) for any choice of $\alpha, \rho \in [0,1]$, $x_{\alpha, \rho}$ is an admissible pure strategy. The proof of (i) is trivial. To prove (ii), we observe that $x_{\alpha, \rho}$ is continuous as a function of t and $x_{\alpha, \rho}(t) \geq 0$ for all $t \in [0,1]$. It remains to show that $\int_0^1 x_{\alpha, \rho}(t) dt = 1$. Thus,

$$\begin{aligned} \int_0^1 x_{\alpha, \rho}(t) dt &= \int_0^1 v(t) dt + \rho \int_0^1 v(t) \cos 2\pi \left(\alpha - \int_0^t v(\lambda) d\lambda \right) dt \\ &= 1 + \rho \int_0^1 \cos 2\pi \left(\alpha - \int_0^t v(\lambda) d\lambda \right) d \int_0^t v(\lambda) d\lambda \\ &= 1 + \rho \int_0^1 \cos 2\pi (\alpha - u) du \\ &= 1 + 0 = 1, \text{ as required.} \end{aligned}$$

To prove that the strategy given by (1), (2) and (3) is optimal for Blotto, we must show that

$$(4) \quad \int_0^1 \int_0^1 M(x_{\alpha, \rho}, y) df_1^*(\alpha) df_2^*(\rho) \geq 0 \text{ for all } y \in X.$$

On making the required substitution, we obtain the following expression for the left member of (4):

$$(5) \quad \int_0^t \int_0^t \left\{ \int_0^t v(t) \operatorname{sgn} \left[v(t) \left(1 + \rho \cos 2\pi \left(\alpha - \int_0^t v(\lambda) d\lambda \right) - y(t) \right) \right] dt \right\} \frac{\rho d\rho d\alpha}{\sqrt{1-\rho^2}}$$

On making the transformation $\theta = 2\pi\alpha$, and using the Fubini theorem, one obtains for (5):


$$(6) \quad \frac{1}{2\pi} \int_0^t v \left\{ \int_0^{2\pi} \operatorname{sgn} \left[v(1 + \rho \cos (\theta - \mu)) - y \right] \frac{\rho d\rho d\theta}{\sqrt{1-\rho^2}} \right\} dt,$$

where, for brevity, we have written v, μ, y , respectively, for $v(t), 2\pi \int_0^t v(\lambda) d\lambda, y(t)$.

Now observe that $\frac{\rho d\rho d\theta}{\sqrt{1-\rho^2}}$ is an element of surface area of a unit hemisphere erected on the polar coordinate plane and centered at the origin, the point on the surface of the hemisphere having cylindrical coordinates (ρ, θ, z) . Moreover, the equation

$$v(1 + \rho \cos (\theta - \mu)) - y = 0$$

is the equation of a straight line in polar coordinates (i.e., a vertical plane in cylindrical coordinates) at a directed distance $\frac{y}{v} - 1$ from the origin, the direction being taken with respect to the ray $\theta = \mu$. (μ can be interpreted as an orientation angle.) Moreover, since $\frac{y}{v} \geq 0$ for all t , it follows that $\frac{y}{v} - 1 \geq -1$. Thus, in evaluating the double integral under the braces of (6), we need consider only two cases. Let $S_1 = \left[t \mid \frac{y}{v} \leq 2 \right]$ and $S_2 = \left[t \mid \frac{y}{v} > 2 \right]$. These cases are illustrated in the accompanying figure.



Now, for $t \in S$, the double integral is evaluated as follows:

The signum function is $+1$ to the right of the line (i.e., in E^+ as illustrated) and -1 to the left. Thus, the integral in question is equal to the area of that portion of the hemisphere to the right of the line less the area to the left. But from a well-known property of the surface area of the sphere, this latter area is proportional to $\frac{Y}{V}$, i.e., it is equal to $\pi \frac{Y}{V}$.

Thus, for $t \in S_1$,

$$\left\{ \iint \right\} = (2\pi - \pi \frac{Y}{V}) - \pi \frac{Y}{V} = 2\pi(1 - \frac{Y}{V})$$

Now for $t \in S_2$, the integral becomes simply the negative of the surface area of the hemisphere, i.e., -2π . But for $t \in S_2$, $\frac{Y}{V} > 2$, whence $-2\pi > 2\pi(1 - \frac{Y}{V})$. Thus, for $t \in S_2$,

$$-2\pi = \left\{ \iint \right\} > 2\pi(1 - \frac{Y}{V}).$$

Hence, in any case, we have

$$(7) \quad \left\{ \iint \right\} \geq 2\pi(1 - \frac{Y}{V}).$$

Now, since v is positive, we may multiply the inequality (7) by $\frac{v}{2\pi}$ and integrate:

$$\begin{aligned}
 (8) \quad \frac{1}{2\pi} \int_0^1 \int_0^1 \{ \int_0^1 \int_0^1 \} dt &\geq \int_0^1 v \left(1 - \frac{y}{v} \right) dt &= \int_0^1 v dt - \int_0^1 y dt \\
 &= 1 - 1 \\
 &= 0.
 \end{aligned}$$

Returning to (6) we see that the left member of (8) is Blotto's expectation. This completes the verification.

References

- [1] O. Gross and R. Wagner: A Continuous Colonel Blotto Game, RAND RM-408.
- [2] O. Gross: The Symmetric Blotto Game, RAND RM-424.